



First evidence of microplastic ingestion by fishes from the Amazon River estuary



Tamyris de Souza e Silva Pegado^{a,*}, Kurt Schmid^a, Kirk O. Winemiller^b, David Chelazzi^c,
Alessandra Cincinelli^{c,d}, Luigi Dei^{c,d}, Tommaso Giarrizzo^a

^a Laboratório de Biologia Pesqueira e Manejo dos Recursos Aquáticos, Universidade Federal do Pará, Av. Perimetral 2651, Terra Firme, 66077830 Belém, Pará, Brazil

^b Department of Wildlife and Fisheries Sciences, Texas A&M University, College Station, TX 77843-2258, USA

^c Consorzio Interuniversitario per lo Sviluppo dei Sistemi a Grande Interfase (CSGI), Via della Lastruccia, 3, 50019 Sesto Fiorentino, Florence, Italy

^d Department of Chemistry “Ugo Schiff”, University of Florence, Via della Lastruccia, 3, 50019 Sesto Fiorentino, Florence, Italy

ARTICLE INFO

Keywords:

Brazil
Bycatch
Pollution
Trophic level

ABSTRACT

This study investigated occurrence of microplastic particles in digestive tracts of fishes from the Amazon River estuary. A total of 189 fish specimens representing 46 species from 22 families was sampled from bycatch of the shrimp fishery. Microplastic particles removed from fish gastrointestinal tracts were identified using Attenuated Total Reflectance – Fourier Transform Infrared (ATR-FTIR). In total, 228 microplastic particles were removed from gastrointestinal tracts of 26 specimens representing 14 species (30% of those examined). Microplastic particles were categorized as pellets (97.4%), sheets (1.3%), fragments (0.4%) and threads (0.9%), with size ranging from 0.38 to 4.16 mm. There was a positive correlation between fish standard length and number of particles found in gastrointestinal tracts. The main polymers identified by ATR-FTIR were polyamide, rayon and polyethylene. These findings provide the first evidence of microplastic contamination of biota from the Amazon estuary and northern coast of Brazil.

1. Introduction

During recent decades, changes in manufacturing and consumer behavior together with insufficient waste management have resulted in accumulation of plastic debris in oceans throughout the world (e.g., Costa and Barletta, 2015; Jambeck et al., 2015), with plastic now composing between 60% and 80% of all marine debris (Barnes et al., 2009). It has been estimated that nearly half of all plastic products are discarded in < 12 months after production (Hopewell et al., 2009). Once introduced into marine ecosystems, plastic waste becomes fragmented as it disperses via wind and oceanic currents (Barnes et al., 2009; Lebreton et al., 2012) and is distributed throughout the water column (Bellas et al., 2016). Plastic debris accumulates not only in the open ocean, but also on beaches, mangrove forests and other coastal habitats (Ivar do Sul and Costa, 2007). Although some plastic debris is dumped directly into marine waters, rivers accumulate discarded material throughout their watersheds and transport it to the oceans (Lechner et al., 2014; Vendel et al., 2017). Unfortunately, rivers and estuaries have received relatively little attention with regard to the plastic pollution problem (Costa and Barletta, 2015), especially within the southern hemisphere (Cannon et al., 2016).

Reports of interactions between marine fauna and plastic debris have increased by 75% over the last two decades, including 267 species reported in 1997 (Laist, 1997) and 693 species reported in 2015 (Gall and Thompson, 2015). Plastic waste in the environment negatively impacts biota, including entanglement of animals within large items (macroplastics) and ingestion of microplastics (particles < 5 mm) by organisms, with subsequent transfer within the food web (Fossi et al., 2012; Cole et al., 2013; Ivar do Sul and Costa, 2014). Ingestion of plastic can affect organisms both physically and physiologically, including direct mortality from entanglement and choking as well as sub-lethal effects, such as compromised feeding, digestion, and reproduction activities (Gregory, 2009; Vendel et al., 2017). Exposure to chemical pollutants that bind to plastic particles has become a major concern, especially when chemicals bioaccumulate in fish destined for human consumption (Teuten et al., 2009). The effects of human consumption of organisms that contain microplastics are still poorly understood. Some evidence has been reported that plastic particles may cause immunotoxic responses, resulting either from chemical exposure or particle-induced mechanical stress (Seltenrich, 2015).

In aquatic and marine environments, plastics undergo a continuous process of disintegration from the action of water and wind causing

* Corresponding author.

E-mail address: t.pegado2@gmail.com (T.d.S.e.S. Pegado).

<https://doi.org/10.1016/j.marpolbul.2018.06.035>

Received 14 March 2018; Received in revised form 9 June 2018; Accepted 11 June 2018

Available online 23 June 2018

0025-326X/ © 2018 Elsevier Ltd. All rights reserved.




Fig. 1. Location of the Amazon River estuary in northeastern Brazil (inset) showing the survey area (yellow shaded area). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

abrasion from contact with solid particles, and through chemical decomposition by exposure to solar radiation (Moore, 2008; Barnes et al., 2009). Plastic debris is classified as macroplastics (particle diameter > 25 mm), microplastics (diameter < 5 mm) (GESAMP, 2015) or mesoplastics (5–25 mm) (Jabeen et al., 2017). Microplastics are further classified according to their origin. Primary microplastics are resin pellets and microbeads used in cleaning products, cosmetics, medicines and other products; secondary microplastics are formed from the fragmentation of larger meso- and macroplastics (Cole et al., 2011). Plastic pellets are used worldwide as a raw material in the production of plastic products (Ogata et al., 2009). With exposure to solar radiation, plastic pellets often lose or change their initial white or translucent coloration and many anthropogenic and biogenic chemicals can be adsorbed by their surface (Endo et al., 2005; Miranda and Carvalho-Souza, 2015). Hydrophobic characteristics of plastics allow them to function as vectors for organic contaminants and heavy metals (Colabuono et al., 2010; Holmes et al., 2012).

Many fishes ingest tiny plastic particles either intentionally or accidentally while feeding in the water column or the benthos (Browne et al., 2010). Most investigations of microplastic ingestion by wild fish have been conducted in the northern hemisphere (e.g. Boerger et al., 2010; Phillips and Bonner, 2015), especially in Europe (e.g. Neves et al., 2015; Bellas et al., 2016; McGoran et al., 2017) and North America (e.g. Carson, 2013; Petters and Bratton, 2016). Microplastic ingestion by fishes in the Southern Hemisphere has been documented by studies performed in Africa (e.g. Biginagwa et al., 2016; Naidoo et al., 2016), Australia (e.g. Cannon et al., 2016), Easter Island (e.g. Ory et al., 2017), Indonesia (e.g. Rochman et al., 2015), and South America (Mizraji et al., 2017; Ory et al., 2017). Studies in Brazil have been conducted in the northeastern and southeastern regions (e.g. Possatto et al., 2011; Ferreira et al., 2016; Silva-Cavalcanti et al., 2017), with no investigations as yet for the northern region that includes the Amazon River estuary.

Brazil's northern coastline has low human population density and contains the world's second-longest, continuous area of largely undisturbed mangrove forest (ca. 7000 km²) (Giarrizzo and Krumme, 2008). In 2016, an extensive and biodiverse reef system (~9500 km²) was discovered offshore from the mouth of the Amazon River (Moura

et al., 2016). This discovery, paired with the fact that 20% of Brazil's fisheries landings come from the northern coast (Krumme et al., 2015), lends urgency to the need to improve knowledge about plastic pollution in the region. Based on experiences in estuaries from northeastern Brazil, Costa and Barletta (2015) identified the Amazon River estuary as a priority area for future studies on marine plastic pollution. The goal of our study was to investigate the presence of microplastics ingestion by fishes from the Amazon River estuary on the coast of Brazil. We hypothesized that quantity and size of the ingested particles increases with fish body size, weight and vertical trophic position within the estuarine food web.

2. Material and methods

2.1. Study area

Brazil's North Coast extends over 1400 km along the states of Amapá and Pará, covering an area of approximately 488,000 km² and a variety of ecosystems including mesophotic reefs, islands, tidal flats, and estuaries with extensive mangrove forests (Marceniuk et al., 2013) (Fig. 1). The region's equatorial humid climate (Kottke et al., 2006) has annual rainfall up to 3300 mm and average annual temperatures of 27.5 to 29.5 °C (Pereira et al., 2009).

The region includes the estuary of the world's largest river, the Amazon, with its mean annual discharge of 6.3 trillion m³ of freshwater, 1.2 billion tons of sediments and 290 million tons of solutes that flow onto the continental shelf (Oltman, 1968; Meade, 1985; Nittrouer et al., 1995). The Amazon's freshwater plume can seasonally expand up to 120 km from the river mouth to the open ocean where salinities can close to zero.

The Amazon's massive freshwater discharge affects oceanographic processes, creating dynamic system of currents and tidal fluxes. The large sediment discharge contributes to high primary and secondary productivity, sustaining important artisanal and commercial fisheries (Neiva and Moura, 1977; Wolff et al., 2000). Estuarine fishes and crustaceans have great economic importance and many of them interact with substrates, influencing physical and chemical processes, including nutrient dynamics (Lana et al., 1996).

2.2. Fish sampling

Fish specimens analyzed in this study were obtained from bycatch of the southern brown-shrimp fishery, which is monitored by the Center for Research and Management of Fishing Resources of Brazil's North Coast (CEPNOR) of the Brazilian Ministry of the Environment (MMA). Samples were obtained from 104 bottom trawls carried out between July 2015 and August 2016 at depths varying from 35 to 85 m. Due to ship's limited freezer space, a sample of the most abundant fish species were collected from each haul, frozen immediately, and stored for up to 2 weeks at -20°C in the laboratory prior to processing.

2.3. Sample processing

In the laboratory, the fishes were identified to species level and each specimen was measured for standard length (SL) with a caliper (0.01 cm precision) and weighed with a digital scale (Marte BL3200H; 0.01 g precision). For each species trophic level was assigned according to values provided in FishBase (Froese and Pauly, 2017). To remove the digestive tract, a longitudinal incision was made in the abdominal area using surgical forceps and a scalpel. Each digestive tract (stomach and intestine) was cut longitudinally, and contents were washed into a Petri dish using 70% ethanol.

Stomach and intestine contents were examined under a stereo-microscope (Opton Tim-2b) at $6.5\times$ to $50\times$ magnification. Plastic particles were separated from other ingested particles, counted, classified according to shape and colour, measured (diameter in longest dimension to 0.001 mm precision), and photographed using a ZEISS SteREO Discovery V12 stereo microscope with the Zen software (blue edition, v2.0, Zeiss, Oberkochen, Germany).

Before and after each procedure for each specimen, all work surfaces and instruments were thoroughly cleaned with 70% ethanol and a new, clean pair of latex gloves was worn. To test for potential presence of airborne plastic fibers at the work station, a clean (using 70% ethanol) glass Petri dish that was placed at the work station at the start of each day was examined with the stereo-microscope at the end of the day.

A sample of each microplastic type recorded in the study was randomly selected for polymer identification. Recent studies showed that Fourier Transform Infrared (FTIR) spectroscopy is the most reliable method to identify the composition of maritime plastic debris, using either single-element or Focal Plane Array (FPA) detectors (Srinivasa Reddy et al., 2006; Mecozzi et al., 2016; Cincinelli et al., 2017). In this study, FTIR spectra were collected in Attenuated Total Reflectance (ATR) mode, using a single-element MCT detector, which was deemed as the optimal set up, given the type and morphology of the plastic samples. The ATR-FTIR analysis of the samples was carried out using a Cary 620–670 FTIR microscope, equipped with a GeATR crystal (Agilent Technologies). The spectra were recorded directly on the samples with a spectral resolution of 8 cm^{-1} , acquiring 128 scans for each spectrum in the $4000\text{--}650\text{ cm}^{-1}$ spectral range.

2.4. Statistical analysis

The percentage of frequency of occurrence of microplastics within digestive tracts was calculated using the following formula: $\text{FO}\% = (\text{Ni} / \text{N}) \times 100$, where $\text{FO}\%$ = frequency of occurrence of microplastic particles; Ni = number of gastrointestinal tracts that contained microplastic particles; N = total number of gastrointestinal tracts examined.

Kendall's rank correlation was performed to assess the association between i) number of ingested microplastic particles and fish body size, ii) number of ingested microplastic particles and fish weight, iii) number of ingested microplastic particles and fish trophic level, and iv) the size of microplastic particles and fish body size. Whenever the correlation was significant, a linear regression was performed to derive

an equation predicting the relationship between the response variable and independent variable. Statistical tests only included specimens that had microplastic particles in the gastrointestinal tract, and were performed using the R statistical package (R Core Team, 2017).

3. Results

Overall, 189 fish specimens representing 46 species from 22 families were analyzed (Table 1). Standard length (\pm SD) averaged $24.7 (\pm 13.3)$ cm, varying from 10.2 to 92 cm, and average body weight was $558.3\text{ g } (\pm 1429.2)$ ranging from 28.2 g to 10,000 g. All fishes are carnivorous and the species trophic level (TL), based on values reported in FishBase, ranged from 3.1 to 4.5.

In total, 228 microplastic particles were recovered from gastrointestinal tracts of 26 specimens belonging to 14 species (Table 1), which represented 13.7% of the total abundance and 30.4% of the total species richness in samples. From the total number of microplastic particles, 210 (92.1%) were found in the stomachs of 21 specimens representing 11 species, and 18 particles (7.9%) were recovered from the intestines of 13 specimens representing six species (Table 1). On average, $1.2 (\pm 5.0)$ particles per fish were found. The standard length of fishes with ingested microplastics varied from 16.0 to 57.5 cm (mean $\text{SL} = 32.7\text{ cm } \pm 11.7$), and weight varied between 72.7 and 10,000 g (mean = $1531.9\text{ g } (\pm 3132.7)$). Size, weight and number of microplastic particles found in gastrointestinal tract for each fish are provided as supplementary material (Appendix 1).

The greatest number of ingested microplastics per specimen (50 particles) was recorded for *C. hippos*. This species was responsible for 40.3% of all recovered microplastic particles. The frequency of occurrence (FO%) of microplastics per species, among those with three or more specimens examined, varied between 18.7% (*Cynoscion microlepidotus*) and 100% (*Bagre marinus*, *Caranx hippos*).

A positive correlation was found between fish SL and number of microplastic particles in the gastrointestinal tract (Kendall's $\text{Tau} = 0.41$, $p = 0.003$). Linear regression analysis indicated that each additional centimeter of SL was associated with an additional 0.58 ingested particles ($r^2 = 0.37$; $p = 0.0008$; $y = -10.374 + 0.586x$) (Fig. 2). In contrast, no significant correlation ($p = 0.43$) was found between the quantity of ingested particles and trophic level, between the size microplastic particles and fish SL ($p = 0.48$), or between the quantity of ingested particles and fishes weight ($p = 0.15$).

The size of plastic particles recovered from fish gastrointestinal tracts was always $< 5\text{ mm}$ (range = $0.38\text{--}4.16\text{ mm}$, average (\pm SD) = $1.82 (\pm 0.68)$). Consequently, all particles are classified as microplastics (Arthur et al., 2009). Particles were recorded as four shape categories: pellets (97.4%), sheets (1.3%), fragments (0.4%) and threads (0.9%). Colours were either clear, yellow, orange or blue (Fig. 3).

Microplastic polymers were identified using ATR-FTIR and comparison with reference spectra (Fig. 3). Samples classified as blue thread and yellow fragments had absorption spectra consistent with polyethylene (Fig. 3a, b): $2920 (\nu_{\text{as}}\text{CH}_2)$, $2850 (\nu_{\text{s}}\text{CH}_2)$, 1471 (bending deformation), $1373 (\delta_{\text{s}}\text{CH}_3)$ and 720 cm^{-1} (rocking deformation) (Gulmine et al., 2002). Among blue and yellow fragments, some spectral bands also could be consistent with presence of adsorbed algae: ca. 3350 (N–H, O–H stretching), 1650 (amide I) and 1540 cm^{-1} (amide II) (Kong and Yu, 2007). Among blue fragments, some weak bands and shoulders suggest the presence of oxygenated groups that form during the abiotic oxidation of polyethylene (Gewert et al., 2015): 1715 (C=O stretching ketones, carboxylic acids) and 1738 cm^{-1} (C=O stretching esters) (Gardette et al., 2013). Transparent sheets (Fig. 3c) showed spectral bands characteristic of Rayon at 3330 (O–H stretching), 2901 (CH, CH_2 stretching), 1648 (HOH bending of water), 1445 (H–C–H and H–O–C bending), 1373 and 1320 (H–C–C, H–C–O, and H–O–C bend), 1110 and 1045 (C–C and C–O–C stretching), and 901 cm^{-1} (C–O–C in plane, symmetric) (Kaur et al., 2013; Li-Ling,

Table 1
Summary information for fishes examined for microplastic particles.

Family	Species	Trophic level	N° of fish	N° of fish with microplastic	Frequency of occurrence (%)	N° of microplastic particles in gastrointestinal tract	Average N° of microplastic particles per specimen ^b	N° of fish with microplastic in stomach	N° of microplastic particles in stomach	N° of fish with microplastic in intestine	N° of microplastic particles in intestine
Ariidae	<i>Bagre bagre</i> ^a	4	7	5	71.4	64	12.8	5	62	2	2
	<i>Bagre marinus</i> ^a	3.5	4	4	100	31	7.8	4	31	0	0
Batrachoididae	<i>Notarius grandicassis</i>	4	4	0	0	0	0	0	0	0	0
	<i>Batrachoides surinamensis</i>	3.7	1	0	0	0	0	0	0	0	0
Carangidae	<i>Caranx crysos</i>	4.1	3	0	0	0	0	0	0	0	0
	<i>Caranx hippos</i> ^a	3.6	3	3	100	92	30.7	3	92	0	0
	<i>Selene setapinnis</i>	3.7	1	0	0	0	0	0	0	0	0
	<i>Selene vomer</i> ^a	4.3	2	1	50	2	2.0	1	2	0	0
	<i>Chaetodipterus faber</i>	4.5	5	0	0	0	0	0	0	0	0
Ephippidae	<i>Anisotremus surinamensis</i>	3.6	1	0	0	0	0	0	0	0	0
	<i>Anisotremus virginicus</i>	3.6	1	0	0	0	0	0	0	0	0
Haemulidae	<i>Conodon nobilis</i>	3.6	8	0	0	0	0	0	0	0	0
	<i>Genyaremus luteus</i>	3.5	8	0	0	0	0	0	0	0	0
	<i>Haemulon plumieri</i>	3.8	13	0	0	0	0	0	0	0	0
	<i>Haemulon steindachneri</i>	3.7	5	0	0	0	0	0	0	0	0
	<i>Orthopristis ruber</i>	3.6	2	0	0	0	0	0	0	0	0
	<i>Lutjanus analis</i> ^a	3.9	3	1	33.3	1	1.0	0	0	1	1
	<i>Lutjanus synagris</i> ^a	3.8	2	1	50	1	1.0	0	0	2	1
	<i>Cynoponiticus savanna</i>	3.5	1	0	0	0	0	0	0	0	0
	<i>Gymnothorax ocellatus</i>	4.1	1	0	0	0	0	0	0	0	0
	<i>Rhinoptera bonasus</i>	3.2	1	0	0	0	0	0	0	0	0
Narcinidae	<i>Narcine brasiliensis</i> ^a	3.2	6	1	16.7	3	3.0	0	0	1	3
	<i>Ophichthus cylindroides</i>	4	1	0	0	0	0	0	0	0	0
Polynemidae	<i>Ophichthus ophis</i>	4.5	1	0	0	0	0	0	0	0	0
	<i>Polydactylus oligodon</i> ^a	3.7	1	1	100	3	3.0	1	3	0	0
	<i>Polydactylus virginicus</i>	3.7	13	0	0	0	0	0	0	0	0
	<i>Pomatomus saltatrix</i>	4.5	2	0	0	0	0	0	0	0	0
Rachycentridae	<i>Rachycentron canadum</i>	4	1	0	0	0	0	0	0	0	0
	<i>Bairdiella ronchus</i>	3.5	4	0	0	0	0	0	0	0	0
Sciaenidae	<i>Ctenosciaena gracilicirrius</i>	3.9	11	0	0	0	0	0	0	0	0
	<i>Cynoscion jamaicensis</i>	3.8	3	0	0	0	0	0	0	0	0
	<i>Cynoscion leiarchus</i> ^a	3.1	2	1	50	2	2.0	1	2	0	0
	<i>Cynoscion microlepidotus</i> ^a	4	16	3	18.7	4	1.3	3	3	1	1
	<i>Cynoscion virescens</i> ^a	4	7	1	14.3	3	3.0	1	3	0	0
	<i>Macrondon ancyodon</i> ^a	3.9	13	1	7.7	2	2.0	1	2	0	0
	<i>Menticirrhus americanus</i>	3.5	1	0	0	0	0	0	0	0	0
	<i>Microgogonias furnieri</i>	3.1	6	0	0	0	0	0	0	0	0
	<i>Paralichthys brasiliensis</i>	3.4	6	0	0	0	0	0	0	0	0
	<i>Scomberomorus brasiliensis</i>	3.3	1	0	0	0	0	0	0	0	0
Scombridae	<i>Epinephelus itajara</i>	4.1	2	0	0	0	0	0	0	0	0
	<i>Sphyrna tiburo</i> ^a	3.9	2	2	100	18	9.0	1	8	2	10
Serranidae	<i>Peprilus paru</i>	4.5	2	0	0	0	0	0	0	0	0
	<i>Stromateidae</i>	3.6	2	0	0	0	0	0	0	0	0
Tetraodontidae	<i>Colomesus psittacus</i>	3.6	2	0	0	0	0	0	0	0	0
	<i>Mustelus canis</i>	3.6	2	0	0	0	0	0	0	0	0
Triakidae	<i>Mustelus higniani</i>	3.6	3	0	0	0	0	0	0	0	0
	<i>Trichiurus lepturus</i> ^a	4.4	5	1	20	2	2.0	1	2	0	0
Trichiuridae			189	26	-	228	1.75	22	210	9	18
Total											

Species in bold font support important commercial fisheries.

^a Fishes that ingested microplastic.

^b Sample based only on specimens that ingested microplastic.




Fig. 2. Scatter plot showing the relationship between the number of ingested microplastic particles and SL among fishes that had microplastic particles in their gastrointestinal tracts (n = 26). Shaded area indicates 95% confidence interval for the linear regression.

2007). Finally, samples classified as pellet I and pellet II (Fig. 3d, e) had spectra indicative of nylon at 3285 (N–H stretching), 3060 (C–H stretching, asym., N–H band), 2953 (CH₂ asym. stretching), 1640 (amide I), 1529 (amide II), 1447 and 1400 (CH₂ bending), and 1242 cm⁻¹ (amide III) (Mahdi, 2011; Charles et al., 2009; Fayemi et al., 2016). For the pellet I sample, the broad band around 1045 cm⁻¹ (C–O–C stretching) suggests the presence of adsorbed cellulosic fibers.

4. Discussion

This study provides the first evidence of ingestion of microplastic particles by fishes from the Amazon River estuary. In recent studies conducted in the northern Atlantic Ocean, 19.80% (Neves et al., 2015), 17.50% (Bellas et al., 2016) and 2.98% (Brate et al., 2016) of fishes had

microplastics in their gastrointestinal tracts. Microplastics were reported from guts of 50% of marine fishes near the city of Salvador (Miranda and Carvalho-Souza, 2015) and in 23% of fishes from the Goiana estuary (Possatto et al., 2011), both in Northeast Brazil. Another study of fishes from the Goiana estuary (Ramos et al., 2012) found levels of microplastic ingestion (13.4%) similar to the level we found for fishes the Amazon estuary (13.7%).

The greatest number of ingested microplastics (50 particles) was encountered in a Crevalle jack (*Caranx hippos*) a predatory fish that can attain 124 cm SL and 32 kg (Meyer et al., 2001; Froese and Pauly, 2017). Considered generalists, Crevalle jack feed on abundant prey, such as schooling fish and crustaceans (Kwei, 1978; Sancho, 2000). A previous study had confirmed the relationship between the ingestion of microplastics and the feeding strategies of demersal fishes in the Eastern Mediterranean (Anastasopoulou et al., 2013). Romeo et al. (2015) proposed that microplastics are ingested most frequently by generalist foragers that target abundant small prey - a characteristic consistent with Crevalle jack feeding habits. The authors further proposed that microplastics also are ingested secondarily when their prey already contain microplastics. Given that microplastics tend to accumulate in the benthos (Bellas et al., 2016), biomagnification may be accelerated in predators that target bottom-feeding fishes and crustaceans. Ingestion of microplastics by the Crevalle jack is cause for concern, given the species' ecological importance in many tropical and subtropical marine ecosystems. Additionally, fishes of the Carangidae family are among the most important in tropical fisheries (Reuben et al., 1992; Crabtree et al., 2002). Our finding of a significant correlation between fish length and the quantity of ingested microplastic particles corroborates those of Alomar and Deudero (2017) for goatfish, *Mullus surmuletus*, in the Western Mediterranean. Our failure to find a relationship between the number of ingested microplastics and trophic level may have been influenced by the limited range of trophic levels (3.1 to 4.5) among the species surveyed.

Several investigations of marine vertebrates and invertebrates have found that microfibrers are the most commonly ingested microplastic particles (Cole et al., 2013; Bellas et al., 2016; Mizraji et al., 2017). In our study, plastic pellets comprised > 97% of the microplastics recovered from fish gastrointestinal tracts. This corroborates the findings





Fig. 3. ATR-FTIR spectra of the microplastic particles: (a) blue thread (polyethylene); (b) yellow fragment (polyethylene); (c) transparent sheet (rayon); (d) pellet I (polyamide); (e) pellet II (polyamide). Highlighted in red the FTIR inset showing the oxidation bands of polyethylene in spectra a and b. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



of Miranda and Carvalho-Souza (2015) that fishes from the Brazil's northeastern coast near the city of Salvador had exclusively ingested pellets. This region is home to the largest petrochemical complex in Latin America, which could contribute to plastic pollution of rivers and coastal habitats (Ogata et al., 2009).

The colour of the recovered pellets (predominately shades of yellow) was similar to that observed by Miranda and Carvalho-Souza (2015) and indicative of persistent organic pollutants (POPs) attached to the surface of plastics (Endo et al., 2005). Furthermore, the yellowish colour suggests that the pellets experienced oxidation and photo-oxidation while adrift in the sea (Ogata et al., 2009; Gewert et al., 2015), implying the original introduction into the environment might have occurred far from our study area. Yellow colour also can be caused by interaction with digestive enzymes (Miranda and Carvalho-Souza, 2015).

As estimated by ATR-FTIR analysis, polyamide (Nylon) represented 97.4% of all microplastics ingested by fishes in our sample from the Amazon estuary. We conclude that the high density (1.13–1.15 g/m³) of polyamide (GESAMP, 2015), paired with the round shape of pellets, contributed to their accessibility to fishes in the study area. Most fishing gear is manufactured from nylon (Timmers et al., 2005), and the fishing industry has been estimated to contribute approximately 18% of all plastic debris found in the oceans (Andrady, 2011).

Apart from the nylon pellets, blue thread, yellow fragments and transparent sheets each contributed < 2% to the total of plastic particles recovered from fish gastrointestinal tracts. Blue thread and yellow fragments were identified as polyethylene (PE), a commonly produced polymer (Jambeck et al., 2015) used to make plastic bags and storage containers (GESAMP, 2015). Globally, PE is the most abundant polymer found in the environment (Andrady and Neal, 2009; Hidalgo-Ruz et al., 2012). Relative to nylon pellets, PE microplastics were uncommon in fish gastrointestinal tracts, and this could be related to their relatively low density (0.91–0.96 g/cm³) (Coutinho et al., 2003; GESAMP, 2015). Floating PE particles, therefore, would be less available to demersal fishes. Transparent sheets were made of rayon, a man-made, semi-synthetic polymer that is used as an artificial textile material (Kauffman, 1993; Woodall et al., 2014). Possible sources are clothing, furniture, and personal hygiene products in sewage effluents (Lusher et al., 2012). In the Amazon estuary, this rayon contributed insignificantly, probably due to this material's rapid rate of degradation (Park et al., 2004).

Some marine systems, such as enclosed bodies (e.g., Mediterranean Sea) and zones of gyre convergence, appear to accumulate large amounts of microplastics (Barnes et al., 2009; Ryan et al., 2009), and rivers are considered major contributors to pollution in these regions (Jambeck et al., 2015). Schmidt et al. (2017) demonstrated that rivers transport microplastics more efficiently than macroplastics, and that the concentration of microplastic particles is positively correlated to river size. Along with mismanagement of solid waste disposal, wastewater discharge, inland navigation and industrial pollution all contribute to entry of microplastics into fluvial ecosystems (Lechner et al., 2014). The Amazon River discharges the world's greatest volume of freshwater. Lebreton et al. (2017) estimated that 38,900 tons of plastic wastes are transported from the Amazon River into the Atlantic Ocean annually, suggesting that most of the microplastics found in our study could have originated in the river.

Microplastics, more than macroplastics, are influenced by advection and circulation patterns, which contributed to the accumulation of particles in deep-sea environments (Woodall et al., 2014; Van Cauwenberghe et al., 2015). Offshore convection, saline subduction, and other oceanographic processes along the coast near the Amazon River Mouth could contribute to accumulate microplastics in marine sediments (Talley et al., 2002; Stabholz et al., 2013).

The United Nations Environment Program (UNEP) listed plastic pollution as a critical problem, comparable to climate change (UNEP, 2014). Research that identifies and quantifies microplastics within the

environment and bodies of organisms will help us to understand the magnitude and scope of microplastic pollution, and pave the way to manage this problem (Boerger et al., 2010; Alomar and Deudero, 2017). Our findings indicate significant ingestion of microplastics by diverse fishes inhabiting the Amazon estuary and half of the species, including many that support important fisheries (Table 1). This raises human health concerns, because ingestion of fish that consume plastics has the potential to increase the body burden of hazardous chemicals that adsorb to plastics in the environment (Rochman et al., 2015) and subsequently bioaccumulate (Oehlmann et al., 2009; Rochman et al., 2013). The degree to which microplastics and associated compounds bioaccumulate in food chains is poorly understood at present.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.marpolbul.2018.06.035>.

Acknowledgments

The authors are grateful to A. Klautau and J. Romão Jr. (CEPNOR) Center for Research and Management of Fishing Resources of Brazil's North Coast of the Brazilian Ministry of the Environment (MMA) in partnership with the Aquatic Ecology Group of the Federal University of Pará (UFPA), for their logistical support during the offshore sampling campaigns in the "Shrimp NEN" research project (#445766/2015-8). We also like to thank the students of the Aquatic Ecology Group (GEA) at the Federal University of Pará (UFPA) for their assistance during the laboratory processing and analysis. We thank the editor and the two referees for their valuable comments on the manuscript. The first (TP) and second (KS) authors were funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). The third author (KW) received support from the International Sportfish Fund and National Science Foundation grant DEB 1257813. The fourth author (DC) and fifth author (AC) gratefully acknowledge CSGI (Consorzio Interuniversitario per lo Sviluppo dei Sistemi a Grande Interfase), Florence (Italy) for financial support. The seventh author (TG) received a productivity grant (CNPq #310299/2016-0) and funding from Fundação de Apoio à Pesquisa, Extensão e Ensino em Ciências Agrárias (FUNPEA) with the project "Estudos Ecológicos e Ecotoxicológicos em Peixes da Região Norte e Nordeste do Brasil".

References

- Alomar, C., Deudero, S., 2017. Evidence of microplastic ingestion in the shark *Galeus melastomus* Rafinesque, 1810 in the continental shelf off the western Mediterranean Sea. *Environ. Pollut.* 223, 223–229. <http://dx.doi.org/10.1016/j.envpol.2017.01.015>.
- Anastasopoulou, A., Mytilineou, C., Smith, C.J., Papadopoulou, K.P., 2013. Plastic debris ingested by deep-water fish of the Ionian Sea (Eastern Mediterranean). *Deep Sea Res., Part I* 74, 11–13. <http://dx.doi.org/10.1016/j.dsr.2012.12.008>.
- Andrady, A.L., 2011. Microplastics in the marine environment. *Mar. Pollut. Bull.* 62, 1596–1605. <http://dx.doi.org/10.1016/j.marpolbul.2011.05.030>.
- Andrady, A.L., Neal, M.A., 2009. Applications and societal benefits of plastics. *Philos. Trans. R. Soc. B* 364, 1977–1984. <http://dx.doi.org/10.1098/rstb.2008.0304>.
- Arthur, C., Baker, J., Bamford, H., 2009. Proceedings of the International Research Workshop on the Occurrence, Effects and Fate of Microplastic Marine Debris. Sept 9–11, 2008. NOAA Technical Memorandum NOS-OR&R-30. <https://repository.library.noaa.gov/view/noaa/2509>, Accessed date: 15 January 2018.
- Barnes, D.K.A., Galgani, F., Thompson, R.C., Barlaz, M., 2009. Accumulation and fragmentation of plastic debris in global environments. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* 364, 1985–1998. <http://dx.doi.org/10.1098/rstb.2008.0205>.
- Bellas, J., Martínez-Armental, J., Martínez-Cámara, A., Besada, V., Martínez-Gomes, C., 2016. Ingestion of microplastics by demersal fish from the Spanish Atlantic and Mediterranean coasts. *Mar. Pollut. Bull.* 109, 55–60. <http://dx.doi.org/10.1016/j.marpolbul.2016.06.026>.
- Biginagwa, F.J., Mayoma, B.S., Shashoua, Y., Syberg, K., Khan, F.R., 2016. First evidence of microplastics in the African Great Lakes: recovery from Lake Victoria Nile perch and Nile tilapia. *J. Great Lakes Res.* 42, 146–149. <http://dx.doi.org/10.1016/j.jglr.2015.10.012>.
- Boerger, C.M., Lattin, G.L., Moore, S.L., Moore, C.J., 2010. Plastic ingestion by planktivorous fish in the North Pacific Central Gyre. *Mar. Pollut. Bull.* 60, 2275–2278. <http://dx.doi.org/10.1016/j.marpolbul.2010.08.007>.
- Brate, I.L.N., Eidsvoll, D.P., Steindal, C.C., Thomas, K.V., 2016. Plastic ingestion by Atlantic cod (*Gadus morhua*) from the Norwegian coast. *Mar. Pollut. Bull.* 112, 105–110. <http://dx.doi.org/10.1016/j.marpolbul.2016.08.034>.

- Brown, M.A., Galloway, T.S., Thompson, R.C., 2010. Spatial patterns of plastic debris along estuarine shorelines. *Environ. Sci. Technol.* 44, 3404–3409. <http://dx.doi.org/10.1021/es903784e>.
- Cannon, S.M.E., Lavers, J.L., Figueiredo, B., 2016. Plastic ingestion by fish in the Southern Hemisphere: a baseline study and review of methods. *Mar. Pollut. Bull.* 107, 286–291. <http://dx.doi.org/10.1016/j.marpolbul.2016.03.057>.
- Carson, H.S., 2013. The incidence of plastic ingestion by fishes: From the prey's perspective. *Mar. Pollut. Bull.* 74, 170–174. <http://dx.doi.org/10.1016/j.marpolbul.2013.07.008>.
- Charles, J., Ramkumaar, G.R., Azhagiri, S., Gunsakeran, S., 2009. FTIR and thermal studies on Nylon-66 and 30% glass fibre reinforced nylon-66. *E-J. Chem.* 6, 23–33. <http://dx.doi.org/10.1155/2009/909017>.
- Cincinelli, A., Scopetani, C., Chelazzi, D., Lombardini, E., Martellini, T., Katsoyannis, A., Fossi, M.C., Corsolini, S., 2017. Microplastic in the surface waters of the Ross Sea (Antarctica): occurrence, distribution and characterization by FTIR. *Chemosphere* 175, 391–400. <http://dx.doi.org/10.1016/j.chemosphere.2017.02.024>.
- Colabuono, F.E., Taniguchi, S., Montone, R.C., 2010. Polychlorinated biphenyls and organochlorine pesticides in plastics ingested by seabirds. *Mar. Pollut. Bull.* 60, 630–634. <http://dx.doi.org/10.1016/j.marpolbul.2010.01.018>.
- Cole, M., Lindeque, P., Halsband, C., Galloway, T.S., 2011. Microplastics as contaminants in the marine environment: a review. *Mar. Pollut. Bull.* 62, 2588–2597. <http://dx.doi.org/10.1016/j.marpolbul.2011.09.025>.
- Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., Galloway, T.S., 2013. Microplastic ingestion by zooplankton. *Environ. Sci. Technol.* 47, 6646–6655. <http://dx.doi.org/10.1021/es400663f>.
- Costa, M.F., Barletta, M., 2015. Microplastics in coastal and marine environments of the Western Tropical and Sub-Tropical Atlantic Ocean. *Environ. Sci.: Processes Impacts* 17, 1868–1879. <http://dx.doi.org/10.1039/C5EM00158G>.
- Coutinho, F.M.B., Mello, I.L., de Santa Maria, L.C., 2003. Polietileno: Principais tipos, propriedades e aplicações. *Polímeros* 13, 1–13.
- Crabtree, R., Hood, P.B., Snodgrass, D., 2002. Age, growth and reproduction of permit (*Trachinotus falcatus*) in Florida waters. *Fish. Bull.* 100, 26–34.
- Endo, S., Takizawa, R., Okuda, K., Takada, H., Chiba, K., Kanehiro, H., Ogi, H., Yamashita, R., Date, T., 2005. Concentration of polychlorinated biphenyls (PCBs) in beached resin pellets: variability among individual particles and regional differences. *Mar. Pollut. Bull.* 50, 1103–1114. <http://dx.doi.org/10.1016/j.marpolbul.2005.04.030>.
- Fayemi, O.E., Adekunle, A.S., Ebenso, E.E., 2016. A sensor for the determination of lindane using PANI/Zn, Fe(III) oxides and nylon 6,6/MWCNT/Zn, Fe(III) oxides nanofibers modified glassy carbon electrode. *J. Nanomater.* 2016, 1:1–10. <http://dx.doi.org/10.1155/2016/4049730>.
- Ferreira, G.V.B., Barletta, M., Lima, A.R.A., Dantas, D.V., Justino, A.K.S., Costa, M.F., 2016. Plastic debris contamination in the life cycle of Acoupa weakfish (*Cynoscion acoupa*) in a tropical estuary. *ICES J. Mar. Sci.* 73, 2695–2707. <http://dx.doi.org/10.1093/icesjms/fsw108>.
- Fossi, M.C., Panti, C., Guerranti, C., Coppola, D., Giannetti, M., Marsili, L., Minutoli, R., 2012. Are baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin whale (*Balaenoptera physalus*). *Mar. Pollut. Bull.* 64, 2374–2379. <http://dx.doi.org/10.1016/j.marpolbul.2012.08.013>.
- Froese, R., Pauly, D. (Eds.), 2017. FishBase. World Wide Web Electronic Publication. www.fishbase.org (version (02/2017)).
- Gall, S., Thompson, R., 2015. The impact of debris on marine life. *Mar. Pollut. Bull.* 92, 170–179. <http://dx.doi.org/10.1016/j.marpolbul.2014.12.041>.
- Gardette, M., Perthue, A., Gardette, J.L., Janecska, T., Földes, E., Pukánszky, B., Therias, S., 2013. Photo- and thermal-oxidation of polyethylene: comparison of mechanisms and influence of unsaturation content. *Polym. Degrad. Stab.* 98, 2383–2390. <http://dx.doi.org/10.1016/j.polymdegradstab.2013.07.017>.
- GESAMP, 2015. Sources, fate and effects of microplastics in the marine environment: a global assessment. Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection. Retrieved from: http://ec.europa.eu/environment/marine/good-environmental-status/descriptor-10/pdf/GESAMP_microplastics%20full%20study.pdf. Accessed, Accessed date: 12 December 2017.
- Gewert, B., Plassmann, M.M., MacLeod, M., 2015. Pathways for degradation of plastic polymers floating in the marine environment. *Environ. Sci.: Processes Impacts* 17, 1513–1521. <http://dx.doi.org/10.1039/c5em00207a>.
- Giarrizzo, T., Krumme, U., 2008. Heterogeneity in intertidal fish fauna assemblages along the world's longest mangrove area in northern Brazil. *J. Fish Biol.* 72, 773–779. <http://dx.doi.org/10.1111/j.1095-8649.2007.01728.x>.
- Gregory, M.R., 2009. Environmental implications of plastic debris in marine setting—entanglement, ingestion, smothering, hangers-on, hitch-hiking and alien invasions. *Philos. Trans. R. Soc.*, B 364, 2013–2025. <http://dx.doi.org/10.1098/rstb.2008.0265>.
- Gulmine, J.V., Janisse, K.P.R., Heise, H.M., Akcelrud, L., 2002. Polyethylene characterization by FTIR. *Polym. Test.* 21 (5), 557–563. [http://dx.doi.org/10.1016/S0142-9418\(01\)00124-6](http://dx.doi.org/10.1016/S0142-9418(01)00124-6).
- Hidalgo-Ruz, V., Gutow, L., Thompson, R.C., Thiel, M., 2012. Microplastics in the marine environment: a review of the methods used for identification and quantification. *Environ. Sci. Technol.* 46, 3060–3075. <http://dx.doi.org/10.1021/es2031505>.
- Holmes, L.A., Turner, A., Thompson, R.C., 2012. Adsorption of trace metals to plastic resin pellets in the marine environment. *Environ. Pollut.* 160, 42–48. <http://dx.doi.org/10.1016/j.envpol.2011.08.052>.
- Hopewell, J., Dvorak, R., Kosior, E., 2009. Plastics recycling: challenges and opportunities. *Philos. Trans. R. Soc.*, B 364, 2115–2126. <http://dx.doi.org/10.1098/rstb.2008.0311>.
- Ivar do Sul, J.A., Costa, D.M.F., 2014. The present and future of microplastic pollution in the marine environment. *Environ. Pollut.* 185, 352–364. <http://dx.doi.org/10.1016/j.envpol.2013.10.036>.
- Ivar do Sul, J.A., Costa, M.F., 2007. Marine debris review for Latin America and the Wider Caribbean Region: from the 1970s until now, and where do we go from here? *Mar. Pollut. Bull.* 54, 1087–1104. <http://dx.doi.org/10.1016/j.marpolbul.2007.05.004>.
- Jabeen, K., Lei, S., Li, J., Dongqi, Y., Tong, C., Jingli, M., Shi, H., 2017. Microplastics and mesoplastics in fish from coastal and fresh waters of China. *Environ. Pollut.* 221, 141–149. <http://dx.doi.org/10.1016/j.envpol.2016.11.055>.
- Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Narayan, R., Law, K.L., 2015. Plastic waste inputs from land into the ocean. *Science* 347, 768–771. <http://dx.doi.org/10.1126/science.1260352>.
- Kauffman, G.B., 1993. Rayon. The first semi-synthetic Fiber product. *Prod. Chem.* 70 (11).
- Kaur, I., Sharma, N., Kumari, V., 2013. Modification of fiber properties through grafting of acrylonitrile to rayon by chemical and radiation methods. *J. Adv. Res.* 4, 547–557. <http://dx.doi.org/10.1016/j.jare.2012.11.003>.
- Kong, J., Yu, S., 2007. Fourier transform infrared spectroscopic analysis of protein secondary structures. *Acta Biochim. Biophys. Sin.* 39, 549–559. <http://dx.doi.org/10.1111/j.1745-7270.2007.00320.x>.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., Rubel, F., 2006. World map of the Köppen-Geiger climate classification updated. *Meteorol. Z.* 15, 259–263. <http://dx.doi.org/10.1127/0941-2948/2006/0130>.
- Krumme, U., Giarrizzo, T., Pereira, R., Silva de Jesus, A.J., Schaub, C., Saint-Paul, U., 2015. Airborne synthetic-aperture radar (SAR) imaging to help assess impacts of stationary fishing gear on the north Brazilian mangrove coast. *ICES J. Mar. Sci.* 72, 939–951. <http://dx.doi.org/10.1093/icesjms/fsu188>.
- Kwei, E., 1978. Food and spawning activity of Caranx hippos off the coast of Ghana. *J. Nat. Hist.* 12, 195–215. <http://dx.doi.org/10.1080/00222937800770081>.
- Laist, D.W., 1997. Impacts of marine debris entanglement of marine life in marine debris including a comprehensive list of species with entanglement and ingestion records. In: Coe, J.M., Rogers, D.B. (Eds.), *Marine Debris—Sources, Impact and Solutions*. Springer-Verlag, New York, pp. 99–139. http://dx.doi.org/10.1007/978-1-4613-8486-1_10.
- Lana, P.C., Camargo, M.G., Brogim, R.A., Isaac, V.J., 1996. Os Bentos da Costa Brasileira: avaliação crítica e levantamento bibliográfico, 1858–1996. *FEMAR, Rio de Janeiro* (431p).
- Lebreton, L.C.M., Van der Zwet, J., Damsteeg, J.W., Slat, B., Andrady, A., Reisser, J., 2017. River plastic emissions to the world's oceans. *Nat. Commun.* 8, 15611. <http://dx.doi.org/10.1038/ncomms15611>.
- Lebreton, L.C.M., Greer, S.D., Borrero, J.C., 2012. Numerical modelling of floating debris in the world's oceans. *Mar. Pollut. Bull.* 64, 653–661. <http://dx.doi.org/10.1016/j.marpolbul.2011.10.027>.
- Lechner, A., Keckeis, H., Lumesberger-Loisl, F., Zens, B., Krusch, R., Tritthart, M., Glas, M., Schludermann, E., 2014. The Danube so colourful: a potpourri of plastic litter outnumbering fish larvae in Europe's second largest river. *Environ. Pollut.* 188, 177–181. <http://dx.doi.org/10.1016/j.envpol.2014.02.006>.
- Li-Ling, C., 2007. Identification of textile fiber by Raman microspectroscopy. *Forensic Sci. J.* 6, 55–62. Available online at: fsjournal.cpu.edu.tw.
- Lusher, A.L., McHugh, M., Thompson, R.C., 2012. Occurrence of microplastics in the gastrointestinal tract of pelagic and demersal fish from the English Channel. *Mar. Pollut. Bull.* 67, 94–99. <http://dx.doi.org/10.1016/j.marpolbul.2012.11.028>.
- Mahdi, H.A., 2011. An FTIR study of characterization of neat and UV stabilized nylon 6,6 polymer films. *J. Pure Appl. Sci.* 24.
- Marceniuk, A.P., Caires, R.A., Wosiacki, W.B., Di Dario, F., 2013. Knowledge and conservation of the marine and estuarine fishes (Chondrichthyes and Teleostei) of the north coast of Brazil. *Biota Neotropica* 13, 251–259. <http://www.biotaneotropica.org.br/v13n4/en/abstract?inventory+bn02613042013>.
- McGoran, A.R., Clark, P.F., Morrirt, D., 2017. Presence of microplastic in the digestive tracts of European flounder, *Platichthys flesus*, and European smelt, *Osmerus eperlanus*, from the River Thames. *Environ. Pollut.* 220, 744–751. <http://dx.doi.org/10.1016/j.envpol.2016.09.078>.
- Meade, R.H., 1985. Suspended sediment in the Amazon River and its tributaries in Brazil during 1982–1984. *U.S. Geol. Surv., Open-File Rep.* 85–492. <https://pubs.usgs.gov/of/1985/0492/report.pdf>.
- Mecozzi, M., Pietroletti, M., Monakhova, Y.B., 2016. FTIR spectroscopy supported by statistical techniques for the structural characterization of plastic debris in the marine environment: application to monitoring studies. *Mar. Pollut. Bull.* 106, 155–161. <http://dx.doi.org/10.1016/j.marpolbul.2016.03.012>.
- Meyer, C.G., Holland, K.N., Wetherbee, B.M., Lowe, C.G., 2001. Diet, resource partitioning and gear vulnerability of Hawaiian jacks captured in fishing tournaments. *Fish. Res.* 53, 105–113. [http://dx.doi.org/10.1016/S0165-7836\(00\)00285-X](http://dx.doi.org/10.1016/S0165-7836(00)00285-X).
- Miranda, D.A., Carvalho-Souza, G.F., 2015. Are we eating plastic-ingesting fish? *Mar. Pollut. Bull.* 103, 109–114. <http://dx.doi.org/10.1016/j.marpolbul.2015.12.035>.
- Mizraji, R., Ahrendt, C., Pérez-Venegas, D., Vargas, J., Pulgar, J., Aldana, M., Ojeda, F.P., Duarte, C., Galbán-Malagón, C., 2017. Is the feeding type related with content of microplastics in intertidal fish gut? *Mar. Pollut. Bull.* 116, 498–500. <http://dx.doi.org/10.1016/j.marpolbul.2017.01.008>.
- Moore, C.J., 2008. Synthetic polymers in the marine environment: a rapidly increasing long-term threat. *Environ. Res.* 108, 131–139. <https://doi.org/10.1016/j.envres.2008.07.025>.
- Moura, R.L., Amado-Filho, G.M., Moraes, F.C., Brasileiro, P.S., Salomon, P.S., Mahiques, M.M., Bastos, A.C., Almeida, M.G., Silva-Jr, J.M., Araújo, B.F., Brito, F.P., Oliveira, B.C.V., Bahia, R.G., Paranhos, R.P., Dias, R.J.S., Siegle, R., Figueiredo-Jr, A.B.G., Pereira, R.C., Leal, C.V., Hajdu, E., Asp, N.E., Gregoracci, G.B., Neumann-Leitão, S., Yager, P.L., Francini-Filho, R.B., Fróes, A., Campeão, M., Silva, B.S., Moreira, A.P.B., Oliveira, L., Soares, A.C., Araújo, L., Oliveira, N.L., Teixeira, J.B., Valle, R.A.B., Thompson, C.C., Rezende, C.E., Thompson, F.L., 2016. An extensive reef system at the Amazon River mouth. *Sci. Adv.* 2, e1501252. <http://dx.doi.org/10.1126/sciadv.2016.09.078>.

- 1501252.
- Naidoo, T., Glassom, D., Smit, A.J., 2016. Plastic ingestion by estuarine mullet, *Mugil cephalus* (L. 1758), in an urban harbour, KwaZulu-Natal, South Africa. *Afr. J. Mar. Sci.* 38, 145–149. <http://dx.doi.org/10.2989/1814232X.2016.1159616>.
- Neiva, G.S., Moura, S.J.C., 1977. Sumário sobre exploração de recursos marinhos do litoral brasileiro: situação atual e perspectivas. Programa de Pesquisa e Desenvolvimento Pesqueiro do Brasil. Série Documentos Ocasionais, Brasília 27. pp. 1–48.
- Neves, D., Sobral, P., Ferreira, J.L., Pereira, T., 2015. Ingestion of microplastics by commercial fish off the Portuguese coast. *Mar. Pollut. Bull.* 101, 119–126. <http://dx.doi.org/10.1016/j.marpolbul.2015.11.008>.
- Nittrouer, C.A., Brunskill, G.J., Figueiredo, A.G., 1995. Importance of tropical coastal environments. *Geo-Mar. Lett.* 15, 121–126. <http://dx.doi.org/10.1007/BF01204452>.
- Oehlmann, J., Schulte-Oehlmann, U., Kloas, W., Jagnytsch, O., Lutz, I., Kusk, K.O., Wollenberger, L., Santos, E.M., Paull, G.C., Van Look, K.J.W., Tyler, C.R., 2009. A critical analysis of the biological impacts of plasticizers on wildlife. *Philos. Trans. R. Soc.*, B 364, 2047–2062. <http://dx.doi.org/10.1098/rstb.2008.0242>.
- Ogata, Y., Takada, H., Mizukawa, K., Hirai, H., Iwasa, S., Endo, S., Mato, Y., Saha, M., Okuda, K., Nakashima, A., Murakami, M., Zurcher, N., Booyatumanondo, R., Zakaria, M.P., Dung, L.Q., Gordon, M., Miguez, C., Suzuki, S., Moore, C., Karapanagiotik, H.K., Weerts, S., McClurg, T., Burres, E., Smith, W., Van Velkenburg, M., Lang, J.S., Lang, R.C., Laursen, D., Danner, B., Stewardson, N., Thompson, R.C., 2009. International PelletWatch: global monitoring of persistent organic pollutants (POPs) in coastal waters. 1. Initial phase data on PCBs, DDTs, and HCHs. *Mar. Pollut. Bull.* 58, 1437–1446. <http://dx.doi.org/10.1016/j.marpolbul.2009.06.014>.
- Oltman, R.E., 1968. Reconnaissance Investigation of Discharge and Water Quality of the Amazon River. US Geological Survey Circular 552, Washington DCpp. 16.
- Ory, N.C., Sobral, P., Ferreira, J.L., Thiel, M., 2017. Amberstripe scad *Decapterus muroadsi* (Carangidae) fish ingest blue microplastics resembling their copepod prey along the coast of Rapa Nui (Easter Island) in the South Pacific subtropical gyre. *Sci. Total Environ.* 586, 430–437. <http://dx.doi.org/10.1016/j.scitotenv.2017.01.175>.
- Park, C.H., Kang, Y.K., Im, S.S., 2004. Biodegradability of cellulose fabrics. *J. Appl. Polym. Sci.* 94, 248–253. <https://doi.org/10.1002/app.20879>.
- Pereira, L.C.C., Dias, J.A., Do Carmo, J.A., Polette, M.A., 2009. Zona Costeira Amazônica Brasileira/the Brazilian Amazon Coastal Zone. *J. Coast. Zone Manag.* 9, 3–7. <http://dx.doi.org/10.5894/rgci172>.
- Petters, C.A., Bratton, S.P., 2016. Urbanization is a major influence on microplastic ingestion by sunfish in the Brazos River Basin, Central Texas, USA. *Environ. Pollut.* 210, 380–387. <http://dx.doi.org/10.1016/j.envpol.2016.01.018>.
- Phillips, M.B., Bonner, T., 2015. Occurrence and amount of microplastic ingested by fishes in watersheds of the Gulf of Mexico. *Mar. Pollut. Bull.* 100, 264–269. <http://dx.doi.org/10.1016/j.marpolbul.2015.08.041>.
- Possatto, F.E., Barletta, M., Costa, M.F., Ivar do Sul, J.A., Dantas, D.V., 2011. Plastic debris ingestion by marine catfish: an unexpected fisheries impact. *Mar. Pollut. Bull.* 62, 1098–1102. <http://dx.doi.org/10.1016/j.marpolbul.2011.01.036>.
- R, 2017. A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Ramos, J.A.A., Barletta, M., Costa, M.F., 2012. Ingestion of nylon threads by Gerreidae while using a tropical estuary as foraging grounds. *Aquat. Biol.* 17, 29–34. <http://dx.doi.org/10.3354/ab00461>.
- Reuben, S., Kasim, H.M., Sivakami, S., Radhakrishnan, P.N., Kurup, K.N., Sivasdas, M., Noble, A., Nair, K.V.S., Raju, S.G., 1992. Fishery, biology and stock assessment of Carangid resources from the Indian seas. *Ind. J. Fish.* 39, 195–234. <http://eprints.cmfri.org.in/id/eprint/526>.
- Rochman, C.M., Hoh, E., Kurobe, T., Teh, S.J., 2013. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. *Sci. Rep.* 3 (3263). <http://dx.doi.org/10.1038/srep03263>.
- Rochman, C.M., Tahir, A., Williams, S.L., Baxa, D.V., Lam, R., Miller, J.T., Teh, F.C., Werorilangi, S., Teh, S.L., 2015. Anthropogenic debris in seafood: Plastic debris and fibers from textiles in fish and bivalves sold for human consumption. *Sci. Rep.* 5, 14340. <http://dx.doi.org/10.1038/srep14340>.
- Romeo, T., Battaglia, P., Peda, C., Consoli, P., Andaloro, F., Fossi, M.C., 2015. First evidence of presence of plastic debris in stomach of large pelagic fish in the Mediterranean Sea. *Mar. Pollut. Bull.* 95, 358–361. <http://dx.doi.org/10.1016/j.marpolbul.2015.04.048>.
- Ryan, P.G., Moore, C.J., Van Franeker, J.A., Moloney, C.L., 2009. Monitoring the abundance of plastic debris in the marine environment. *Philos. Trans. R. Soc. B* 364, 1999–2012. <http://dx.doi.org/10.1098/rstb.2008.0207>.
- Sancho, G., 2000. Predatory behaviors of *Caranx melampygus* (Carangidae) feeding on spawning reef fishes: a novel ambushing strategy. *Bull. Mar. Sci.* 66, 487–496. <https://www.researchgate.net/publication/233698439>.
- Schmidt, C., Krauth, T., Wagner, S., 2017. Export of plastic debris by rivers into the sea. *Environ. Sci. Technol.* 51, 12246–12253. <http://dx.doi.org/10.1021/acs.est.7b02368>.
- Seltenrich, N., 2015. New link in the food chain? Marine plastic pollution and seafood safety. *Environ. Health Perspect.* 123, A34. <http://dx.doi.org/10.1289/ehp.123-A34>.
- Silva-Cavalcanti, J.S., Silva, J.D.B., França, E.J., Araújo, M.C.B., Gusmão, F., 2017. Microplastics ingestion by a common tropical freshwater fishing resource. *Environ. Pollut.* 221, 218–226. <http://dx.doi.org/10.1016/j.envpol.2016.11.068>.
- Srinivasa Reddy, M., Basha, S., Adimurthy, S., Ramachandriah, G., 2006. Description of the small plastics fragments in marine sediments along the Alang-Sosiya ship-breaking yard, India. *Estuar. Coast. Shelf Sci.* 68, 656–660. <http://dx.doi.org/10.1016/j.ecss.2006.03.018>.
- Stabholz, M., Durrieu de Madron, X., Canals, M., Khripounoff, A., Taupier-Letage, I., Testor, P., Heussner, S., Kerhervé, P., Delsaut, N., Houper, L., Lastras, G., Dennielou, B., 2013. Impact of open-ocean convection on particle fluxes and sediment dynamics in the deep margin of the Gulf of Lions. *Biogeosciences* 10, 1097–1116. <http://dx.doi.org/10.5194/bg-10-1097-2013>.
- Talley, J.W., Ghosh, U., Tucker, S.G., Furey, J.S., Luthy, R.G., 2002. Particle-scale understanding of the bioavailability of PAHs in sediments. *Environ. Sci. Technol.* 36, 477–483. <http://dx.doi.org/10.1021/es010897f>.
- Teuten, E.L., Saquing, J.M., Knappe, D.R.U., Barlaz, M.A., Jonsson, S., Björn, A., Rowland, S.J., Thompson, R.C., Galloway, T.S., Yamashita, R., Ochi, D., Watanuki, Y., Moore, C., Viet, P.H., Tana, T.S., Prudente, M., Boonyatmanond, R., Zakaria, M.P., Akkhavong, K., Ogata, Y., Hirai, H., Iwasa, S., Mizukawa, K., Hagino, Y., Imamura, A., Saha, M., Takada, H., 2009. Transport and release of chemicals from plastics to the environment and to wildlife. *Philos. Trans. R. Soc. B* 364, 2027–2045. <http://dx.doi.org/10.1098/rstb.2008.0284>.
- Timmers, M.A., Kistner, C.A., Donohue, M.J., 2005. Marine Debris of the Northwestern Hawaiian Islands: Ghost Net Identification. U.S. Dept of Commerce, National Oceanic and Atmospheric Administration, National Sea Grant College Program (31 pages. http://www.nmfs.gov/asset_upload/library/86086/n20864gx_cuwr_conservancy_001_3714).
- UNEP, 2014. UNEP Year Book 2014: Emerging Issues Update. United Nations Environment Programme, Nairobi, Kenya.
- Van Cauwenbergh, L., Claessens, M., Vandegheuchte, M.B., Janssen, C.R., 2015. Microplastics are taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) living in natural habitats. *Environ. Pollut.* 199, 10–17. <http://dx.doi.org/10.1016/j.envpol.2015.01.008>.
- Vendel, A.L., Bessa, F., Alves, V.E.N., Amorim, A.L.A., Patrício, J., Palma, A.R.T., 2017. Widespread microplastic ingestion by fish assemblages in tropical estuaries subjected to anthropogenic pressures. *Mar. Pollut. Bull.* 117, 448–455. <http://dx.doi.org/10.1016/j.marpolbul.2017.01.081>.
- Wolff, M., Koch, V., Isaac, V., 2000. A trophic flow model of the Caeté mangrove estuary (North Brazil) with considerations for the sustainable management of its resources. *Estuar. Coast. Shelf Sci.* 50, 789–803. <http://dx.doi.org/10.1006/ecss.2000.0611>.
- Woodall, L.C., Sanchez-Vidal, A., Canals, M., Paterson, G.L.J., Coppock, R., Sleight, V., Calafat, A., Rogers, A.D., Narayanaswamy, B.E., Thompson, R.C., 2014. The deep sea is a major sink for microplastic debris. *R. Soc. Open Sci.* 1, 140317. <http://dx.doi.org/10.1098/rsos.140317>.